
Flexible - Sigmoidoscopy Essential information for patients





Flexible – Sigmoidoscopy


It is an examination, which allows the doctor to look directly at the lining of the lower part (rectum and sigmoid colon) of the large bowel. It is performed using a thin flexible tube (colonoscope) with a camera and a light, on the tip, which is used to look at the area being examined. The doctor guides the scope through the back passage (anus) and into the colon (large bowel).

It allows us to:

- 1. Take a sample of tissue (biopsy) from the lining of the bowel.
- 2. Remove polyps that can grow on the bowel wall.
- 3. Treat Haemorrhoids (piles) with banding or injection of oily phenol.

The procedure last 10 minutes. It is usually not painful, but can feel a little uncomfortable.

Sigmoidoscopy



There are two options available to you:

1. No sedation option:

- This procedure can be carried out without sedation.
- You will be fully aware of the procedure which most patients find acceptable and not too uncomfortable.
- You will not have to fast and may go home once you have spoken with your doctor.

2. Intravenous sedation option:

If you choose to have conscious sedation you will have a small tube called a cannula put into a vein, usually in your hand or lower arm.

- Conscious sedation will be administered through this cannula.
- It will make you sleepy and relaxed.
- It may make you fall asleep during the procedure, but it is completely normal to be awake. It has an amnesia effect, therefore making you forget.

If you chose to have sedation, you must arrange someone to escort you home and stay with you for 12 hours after the procedure.

He/she should come with you for the appointment or be contactable by phone when you are ready to leave. The nurse will need this person's contact details during admission process.

Patients who have sedation, will be required to stay in the hospital a minimum of 4 hours.

If you do not have an escort or have not arranged for someone to collect you, then your procedure will be cancelled.

Risks of a flexible sigmoidoscopy

Drug Reaction: The sedation used for the procedure is safe with only a very slight risk of reaction.

Bleeding: If a polyp is found during the course of the procedure, this will be removed which can result in some bleeding. This bleeding can often be stopped straight away but may occasionally be more serious or may even occur a few days later. Depending on the size, the risk of bleeding is 1 in 200 polyps removed. Many polyps are the type which can turn into cancer if left untreated for a long period of time and removing them when still benign eliminates the risk.

Perforation: If you do not have polyps removed there is a less than 1 in 2000 chance of causing a perforation, which is a little tear in the lining of the bowel. If polyps are removed, this may occur in 1 in 500 cases. This may extend your hospital stay for a short time and you will need to be given antibiotics, or it may require an operation to repair the tear.

Missed Pathology: It is important to remember that no test is perfect and even with complete examination, a significant disease may be missed in up to 20% of cases.

About the Consent Form

Before a doctor or health care professional examines or treats you, they will need to gain your consent. This will be required in writing. If you later change your mind, you are entitled to withdraw consent even after signing the document.

- Please read the flexible sigmoidoscopy patient information leaflet, prior to signing your consent form.
- Please read the consent form in full, prior to your procedure.
- You will meet the doctor on the day of your procedure, to ask any questions or concerns you may have.

Before your flexible-sigmoidoscopy

Fasting

If you opt for sedation, you must not eat for 6 hours prior to your procedure.

You must not drink for 2 hours prior to your procedure.

If you are NOT having sedation there are no restrictions on eating and drinking.

To obtain a clear view, the lower colon must be completely empty.

There are two ways to clear the bowel for this procedure – Your Doctor will instruct you:

- A) Full Bowel prep drinking a laxative it is essential that you follow these exact instructions.
- **B)** Alternatively, you may be prescribed an enema (a liquid laxative, administered via the back passage) on the day of your procedure. This will be administered by a nurse when you arrive to the unit.

Medicines

You should continue to take your regular medications as normal on the day of your procedure.

- Iron Tablets These will need to be stopped one week before the procedure – as the iron makes your faeces look black in colour, which can be mistaken for blood in the bowel during the examination
- If you are diabetic and you did not receive information from the hospital, please contact the Endoscopy Unit on 01 650 4617
- If you are taking any of these medications

 Aspirin, Warfarin, Clopidrogel (Plavix),
 Rivaroxaban (Xaralto), Dabigatran
 (Pradaxa), Apixaban (Eliquis) phone the
 Endoscopy Unit for advice on 01 650 4617
- If you use a GTN spray for angina please bring this with you.
- If you are asthmatic please bring your inhalers with you.

Please bring a list detailing your regular medications. If you are in doubt, please contact your GP or local pharmacy to provide the list.

Interpretation services

If you require an interpretation service please ask an English-speaking person to call us as soon as possible so that an interpreter can be arranged for you

Special Requirements

If you use a wheelchair or have any physical and/or other disabilities, please contact the unit to arrange any support that you may need to make your endoscopy journey as easy as possible for you.

On Arrival

- On arrival at the hospital, go to the Endoscopy reception area on the second floor.
- The receptionist will take some information regarding your name, address, date of birth and next of kin.
- You will be admitted to the unit by a nurse who will ask information regarding your health history and medical conditions.
- You will be given opportunity to ask any questions or concerns you may have at this stage.
- You will be given a hospital gown and disposable shorts to wear for the procedure.
- You will be asked to wear your dressing gown and slippers, once changed into your gown.
- A small tube called a cannula will be put into a vein. This will happen even if you choose not to have sedation.
- Once the admission assessment is completed by the nurse, you will be brought to a waiting lounge area, where the team from the procedure room will collect you from.

Please bring dressing gown and slippers with you. You are advised not to bring any valuables or any large sums of the money into hospital with you.

During your flexible sigmoidoscopy

- When you are brought into the procedure room, you will meet two registered nurses and the Consultant.
- In the procedure room, you will be made comfortable on the bed, lying on your left side with your knees drawn up.
- You will be able to see the endoscopy screen in front of you.
- You will receive oxygen through your nose.
- A monitor will be placed on your finger to measure your oxygen levels and your blood pressure will be taken every few minutes via a blood pressure cuff on your arm.
- A nurse will look after you throughout the procedure, they will monitor your pulse, blood pressure and heart rate once sedation is given.
- A nurse and doctor will ensure that you are feeling comfortable throughout the procedure.
- You will be given sedation to make you feel relaxed and drowsy but not entirely asleep or unconscious (this is not a general anaesthetic). This will be administered before the procedure starts through the cannula in your arm / hand, and given a few minutes to start working, before the procedure begins.
- The endoscopist will gently pass the scope in through your back passage.
- Air will be passed into the bowel via the scope, this may cause you some discomfort but will pass quickly.
- The procedure takes approximately 15-30 minutes – however can take longer if biopsies need to be taken or polyps need to be removed.

After your flexiblesigmoidoscopy

- You will be brought to the Endoscopy recovery area and your bed.
- Your heart rate and blood pressure will be monitored by a registered nurse.
- Here you can have a rest for a while after your procedure.
- Following your procedure, you may experience some crampy pains and bloating. This is due to trapped wind, it will settle down soon after your procedure. You are encouraged to pass the wind during and after your procedure for your comfort.
- Once you are alert and the sedation has worn off, you will be asked to get up and dressed from the bed.
- You will be escorted by the nurse to a chair in the Endoscopy discharge lounge and handed over to the nurse in this area. Here you will be given some food. You will have a choice of brown or white toast, hot or cold drinks. We do cater for dietary needs, so please let the nurse know when they are ordering your food.
- Your escort should collect you from the Endoscopy Discharge area also located on the 2nd floor. We always wait for your escort to arrive to the unit before giving you any discharge information as you tend to forget it, due to the sedation.
- Your cannula will be removed from you hand or arm once you have had something to eat, passed urine and are not experiencing any side effects from the procedure or medications.
- It is the decision of the consultant performing the test if he/she will be seeing you immediately following the test or if an appointment will be made in the outpatient rooms.
- The consultant's secretary will arrange this appointment for you if needed.

If you have received sedation:

You must not drive, drink alcohol, operate machinery or sign important documents for the next 24 hours.

Please note that following a sedated procedure, you **WILL NOT** be permitted to leave the unit unaccompanied.

Results

- After your procedure, when the sedation has worn off, a doctor/nurse will speak to you about the outcome of the procedure.
- Any biopsies or polyps removed are sent to the laboratory for analysis to assist the doctor in diagnosis.
- We will send a copy of the Endoscopy report to your GP and any lab results should follow within 2-3 weeks.
- The consultant that performed your Colonoscopy may request to see you in their outpatient clinic after the procedure, usually in a couple of weeks and will go through your results.

Any advice or recommended further tests will be given upon discharge.

Frequently Asked Questions

If the symptoms have stopped before my endoscopy, should I still come for the test?

Yes. It is important that you still arrive for your test. Your doctor has organised this test to ensure you have no bowel issues. Although symptoms may have subsided, it is important to proceed in order to ensure all is clear.

Does the procedure hurt?

Generally, no. Sedation is very effective for most people and many people who have sedation do not even remember the procedure.

You may feel cramps or pressure in the back passage during and after the procedure. This is because air is passed through the scope into the bowel to allow the doctor see inside. Every effort is made to minimise this and it usually settles quickly.

The removal of biopsies and polyps does not cause pain.

Can I drive after the procedure?

No, you may not drive for 24 hours post sedation.

The sedation remains in your system for 24 hours. Therefore, it can cause you to feel drowsy at any time within this period. You are not covered by your insurance in the unfortunate event that you are involved in an accident while driving under the influence.

We advise for you not to get public transport as you may feel a little dizzy after the sedation. Escorted home via taxi or your NOK drives you is ideal.

Can I return to work after the procedure?

If you have not had sedation – yes you can return to work

If you were given sedation it will depend on the time of your appointment.

Morning appointments may be okay to return to work the following morning.

If you have an afternoon appointment, it is dependent on how you feel and how much sedation was given on the day of your procedure.

What should I expect when I am sent home?

If sedated, you must be accompanied home by a responsible adult. The sedation used will make you forget things. You may find your memory of the few hours immediately after your flexible sigmoidoscopy to be impaired.

You may experience some abdominal discomfort for 24 hours. If you have had biopsies taken, polys removed or any procedure on haemorrhoids you may experience slight bleeding from the back passage.

Should you experience any sudden onset of pain or sudden blood loss, go to your closest Accident and Emergency Department.

You may resume normal daily activities the day following your sigmoidoscopy.

You will be given discharge information details on the day.

What are the key things to remember?

It's your decision! It's up to you to choose whether or not to consent to what is being proposed. Ask as many questions as you like and please express any concerns you may have about medication, allergies or past medical history

Do I have everything ready for my procedure?

- List of Medication and Medical history.
- Details of your health insurance if applicable.
- Slippers and dressing gown.
- I am fasting from food for six hours before my procedure time, - if I choose to have sedation.
- I am fasting from drinks for 2 hours before my procedure.
- I have arranged an escort to take me home if I choose to have sedation

Location and Parking Information

Beacon Hospital is located in Sandyford, Dublin 18 and can be accessed by Luas, Dublin Bus and car.

LUAS

The Luas (green line) goes directly from St. Stephen's Green to Sandyford. Beacon Hospital, Dublin is located in Sandyford, 10 minutes walk from the Kilmacud Luas station. On exiting the Luas at the Kilmacud station, take the lift/stairs and follow the pathway to the entrance of the station. Turn left here and you will see Beacon Hospital located at the top of the road at the crossroads (this is right beside the Beacon Hotel).

DUBLIN BUS

Beacon Hospital is serviced by the following bus routes:

– No. 11 bus from Wadelai Park (via O'Connell Street). Alight at stop no. 4847 Blackthorn Drive, Luas Car Park.

No. 75 bus from Tallaght/Dun Laoghaire.
 Alight at stop no. 449 Blackthorn Road,
 Corrig Road.

CAR PARKING

An underground public car park is available by turning right on entering the Beacon Court car park.

You may pay your ticket fee on exiting the hospital lobby.





DRIVING

A If coming from the Leopardstown Racecourse direction

- 1. Take the Sandyford Industrial Estate exit.
- 2. Pass 'Woodies' on the left and keep going straight on.
- 3. At the T-junction at the end of this road, turn left (signposted 'Local Traffic').
- 4. The Hospital is at the end of this road on your left on the corner. You will see the entrance to the car park

B If coming from M50, travelling southbound

- 1. Take the Sandyford Industrial Estate exit.
- 2. Pass 'Woodies' on the left and keep going straight on.
- 3. At the T-junction at the end of this road, turn left (signposted 'Local Traffic').
- 4. The Hospital is at the end of this road on your left on the corner. You will see the entrance to the car park here.

C If coming from M50, travelling northbound

- 1. Take the Sandyford Industrial Estate exit.
- 2. Pass 'Woodies' on the left and keep going straight on.
- 3. At the T-junction at the end of this road, turn left (signposted 'Local Traffic').
- 4. The Hospital is at the end of this road on your left on the corner. You will see the entrance to the car park here.

D If coming from Dundrum Village via Upper Kilmacud Road

- 1. Follow the Upper Kilmacud Road to the T junction after the 'Holywell' estate (estate is on your right).
- 2. Take a right at this T-junction.
- 3. Keep straight until you see Beacon Hotel on your left.
- 4. Take a left, and then first right. The entrance to the car park will now be on your right.

Notes / Things to remember

Beacon Hospital Sandyford, Dublin 18 Tel +353 1 293 6600

Endoscopy Unit Opening Hours: Monday - Friday 7:00 - 19:30 hours

Enquiry Lines open Monday to Friday:

Telephone - 01 650 4617 for Clinical Enquires, between 7:30 - 19:30 hours.

Telephone - 01 293 8656 for Scheduling Enquires, between 8:00 - 16:00 hours.



This is Modern Medicine



beaconhospital.ie